Back to school

Low Blood Sugar Readiness Checklist

Equip your child and their school to handle low blood sugar emergencies with confidence

PARTNER WITH YOUR CHILD'S HEALTHCARE PROVIDER

TAKINER WITH TOOK ONLED OTHEREINGARET ROVIDER
Create or update your child's low blood sugar emergency plan.
☐ Discuss if Gvoke HypoPen® is right for your child.
☐ If it is, request a prescription for Gvoke HypoPen 2-Pack™ and the copay card*.
GATHER ESSENTIAL ITEMS
Pack a supply of oral fast-acting carbohydrates, such as glucose tablets, juice, or fruit snacks.
Fill your child's prescription for Gvoke HypoPen and coordinate with the school nurse or administration on its storage and accessibility.
PREPARE YOUR CHILD
Educate your child about the signs and symptoms of low blood sugar.
☐ Stock them up with a supply of "low snacks."
Familiarize them with the location of their emergency supplies and whom to approac for assistance.
☐ Train them on the proper use of Gvoke HypoPen.
INFORM SCHOOL PERSONNEL
Educate teachers, school nurses, and other caregivers about low blood sugar signs and symptoms.
Review the plan so that they know what actions to take if your child experiences a low blood sugar episode.
Communicate the location of your child's Gvoke HypoPen device and provide instructions on when and how to use it.



Scan the QR code to discover more resources at GvokeGlucagon.com



^{*} Offer not valid for prescriptions reimbursed under Medicaid, a Medicare drug benefit plan, TRICARE, or other federal or state health programs (such as medical assistance programs).

INDICATION AND SAFETY SUMMARY

GVOKE is a prescription medicine used to treat very low blood sugar (severe hypoglycemia) in adults and kids with diabetes ages 2 years and above. It is not known if GVOKE is safe and effective in children under 2 years of age.

WARNINGS

Do not use GVOKE if:

- you have a tumor in the gland on top of your kidneys (adrenal gland), called a pheochromocytoma.
- you have a tumor in your pancreas called an insulinoma.
- you are allergic to glucagon or any other inactive ingredient in GVOKE.

GVOKE MAY CAUSE SERIOUS SIDE EFFECTS, INCLUDING:

High blood pressure. GVOKE can cause high blood pressure in certain people with tumors in their adrenal glands.

Low blood sugar. GVOKE can cause low blood sugar in certain people with tumors in their pancreas called insulinomas by making too much insulin in their bodies. Serious allergic reaction. Call your doctor or get medical help right away if you have a serious allergic reaction including:

- rash
- · difficulty breathing
- low blood pressure

COMMON SIDE EFFECTS

The most common side effects of GVOKE in adults include:

- nausea
- vomiting
- · swelling at the injection site
- headache

The most common side effects of GVOKE in children include:

- nausea
- low blood sugar
- · high blood sugar
- vomiting
- abdominal pain
- headache
- · pain or redness at the injection site
- itching

These are not all the possible side effects of GVOKE. For more information, ask your doctor. Call your doctor for medical advice about side effects. You are encouraged to report side effects of prescription drugs to the FDA. Visit fda.gov/medwatch, or call **1-800-FDA-1088**.

BEFORE USING

Before using GVOKE, tell your doctor about all your medical conditions, including if you:

- have adrenal gland problems
- have a tumor in your pancreas
- have not had food or water for a long time (prolonged fasting or starvation)
- have low blood sugar that does not go away (chronic hypoglycemia)
- · are pregnant or plan to become pregnant
- · are breastfeeding or plan to breastfeed
- Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

HOW TO USE

- Read the detailed Instructions for Use that come with GVOKE.
- Use GVOKE exactly how your healthcare provider tells you to use it
- Make sure your relatives, close friends, and caregivers know where you store GVOKE and how to use it the right way before you need their help.
- Act quickly. Having very low blood sugar for a period of time may be harmful.
- Your healthcare provider will tell you how and when to use GVOKE.
- After giving GVOKE, your caregiver should call for emergency medical help right away.
- If you do not respond after 15 minutes, your caregiver may give you another dose, if available. Tell your healthcare provider each time you use GVOKE. Low blood sugar may happen again after receiving an injection of GVOKE. Your diabetes medicine may need to be changed.

HOW TO STORE

- Keep GVOKE in the foil pouch until you are ready to use it.
- Store GVOKE at temperatures between 68°F and 77°F.
- Do not keep it in the refrigerator or let it freeze.

Keep GVOKE and all medicines out of the reach of children.

For more information, call **1-877-937-4737** or go to GvokeGlucagon.com

Please see the Full Prescribing Information for Gvoke

